

IN THE CLAIMS:

Please cancel claims 13 and 14, without prejudice or disclaimer. Please amend the claims pursuant to 37 C.F.R. 1.121 as follows (see the accompanying "marked up" version pursuant to 1.121):

5. (Twice amended) A method for inducing an anti-tumor response in a mammalian patient suffering from a tumor, which method comprises administering to said patient a composition comprising a tumor cell or tumor cell extract with an adjuvant, wherein the tumor cell or tumor cell extract is:

- (i) conjugated to a hapten;
- (ii) of the same tumor type as the patient's tumor;
- (iii) not allogeneic to said patient; and
- (iv) incapable of growing in the body of the patient after injection;

and repeating said administration at weekly intervals,
wherein a therapeutically effective amount of cyclophosphamide is
administered only prior to the first administration of the composition, wherein the
patient is not sensitized to the hapten prior to administration of the composition, and
wherein the composition, when administered with the adjuvant, elicits an anti-tumor
response.

Please add the following new claims:

27. (New) A method for inducing an anti-tumor response in a mammalian patient suffering from a tumor, which method comprises administering to the patient a composition comprising an adjuvant and a tumor cell or tumor cell extract, wherein the tumor cell or tumor cell extract is:

- (i) conjugated to a hapten;
- (ii) of the same tumor type as the patient's tumor;
- (iii) not allogeneic to said patient, and
- (iv) incapable of growing in the body of the patient after injection;

wherein a therapeutically effective amount of cyclophosphamide is administered only prior to the first administration of the composition, wherein the patient is not sensitized to the hapten prior to administration of the composition, and wherein the composition elicits an anti-tumor response.

28. (New) The method of claim 27, which comprises repeating said administration of said composition every week or every two weeks.

29. (New) The method of claim 27, wherein said composition is administered at least three times.

30. (New) The method of claim 27 wherein said therapeutically effective amount of cyclophosphamide comprises about 300 mg/M² of cyclophosphamide, and is administered 3 days prior to administration of the composition.

31. (New) The method of claim 27, wherein said mammalian patient is a human, said tumor cell or extract is a melanoma tumor cell or extract, said hapten is dinitrophenyl, and said adjuvant is *Bacillus Calmette-Guerin*.